

****TITLE****

ASP Conference Series, Vol. ****VOLUME****, ****YEAR OF PUBLICATION****

****NAMES OF EDITORS****

Determining the abundance of extragalactic planets

M. Dominik

*University of St Andrews, School of Physics & Astronomy, North
Haugh, St Andrews, KY16 9SS, United Kingdom*

Abstract. Gravitational microlensing provides a unique technique to reveal information about extragalactic planets. A network of at least four 2m-class telescopes distributed around the northern hemisphere could probe 15–35 jupiters and 4–10 saturns around M31 stars per year.

Planet detection by microlensing

By means of creating distortions to the gravitational field of their parent stars, *unseen planets* of mass m around *unseen lens stars* of mass M can cause deviations of 1–20 % (lasting days for jupiters to hours for earths) to the light curves of background stars undergoing a microlensing event which lasts ~ 1 month. The probability for a planetary signal varies with the projected orbital radius of the planet r_p reaching a maximum near $r_p \sim r_E$, with the *Einstein radius* $r_E = \sqrt{2R_S D}$, the Schwarzschild radius of the lens star $R_S = (2GM)/c^2$, $D = D_L(D_S - D_L)/D_S$, and D_S and D_L being source or lens distances.

By achieving a galactic length scale $D \sim 2.5 \text{ kpc} [r_E/(2.5 \text{ AU})]^2$ for M-dwarf lens stars ($M \sim 0.3 M_\odot$), one becomes sensitive to planetary systems similar to our own. This condition is met not only for observations of Galactic bulge stars being lensed by stars in both the Galactic disk or the bulge itself ($D_S \sim 8.5 \text{ kpc}$, $D_L \sim 6 \text{ kpc}$), but also for observations of stars in other galaxies as M31 (Crotts 1992; Jetzer 1994) lensed by stars in the same galaxy ($D_S \gg D_S - D_L$, $D \approx D_L$).

While the probability to detect a planetary deviation and its duration decreases with smaller mass ratios $q = m/M$ (roughly as $\sqrt{q}$), the maximal deviation δ_{th} is only limited by the radius of the source star $R_\star$ (Bennett & Rhie 1996). For $\delta_{\text{th}} = 10\%$ on observing M31 stars, a minimal mass ratio of $q_{\text{min}} = 2 \times 10^{-3} [R_\star/(100 R_\odot)]^2$ is required for $r_E \sim 2.5 \text{ AU}$ and M-dwarf lens stars, corresponding to planets with twice the mass of Saturn for $R_\star \sim 100 R_\odot$.

M31 microlensing observations

Due to its small angular size, the monitoring of M31 requires only a few fields and therefore only a fraction of telescope time during each night. A proposed network of at least four 2m-class telescopes distributed around the northern hemisphere (Dominik 2002) will provide a sampling interval of $\sim 6 \text{ h}$. *On-line data reduction* based on *difference-imaging* will allow an increased sampling through target-of-opportunity observations in the case of ongoing anomalies.

Since M31 observations comprise *unresolved star fields*, only events with *high peak magnification* A_0 on *bright sources* are detectable, ruling out catching signatures of Earth-mass planets. The smaller fraction of useful events is partly compensated by the *large number of source stars* and an *increased planet detection efficiency* for larger peak magnifications (Griest & Safizadeh 1998).

The table compares the detection capabilities for Galactic and extragalactic planets with ground-based campaigns. Planet detection efficiencies follow Gould & Loeb (1992) and Covone et al. (2000) and are averaged over the lensing zone ($0.6 r_E \leq r_p \leq 1.6 r_E$). The determination of planet parameters is not required for obtaining upper abundance limits based on the absence of deviations.

Table 1. Planet detection capabilities for an M31 network and the PLANET campaign (Dominik et al. 2002) on Galactic bulge stars

	Galactic bulge	M31
number of source stars	$\sim 10^7$	$\sim 10^{10}$
resolution of source stars	resolved	unresolved
telescope time	dedicated	0.5–2.5 h per night
field of view [sq deg]	0.004–0.03	0.01–1
number of fields	~ 20	1–8
monitored during night		
mean sampling interval	1.5–2.5 h	4–6 h
total event rate [yr^{-1}]	~ 300 –600	~ 150 –400
useful types	giants	giants
of source stars	main-sequence stars	
useful peak magnifications	$A_0 \gtrsim 2$	$A_0 \gtrsim 10$
rate of useful events [yr^{-1}]	~ 75	~ 35 –100
planet detection efficiency	$\sim 20\%$ (jupiters) $\sim 1.5\%$ (earths)	$\sim 35\%$ (jupiters) $\sim 10\%$ (saturns)
planet probing rate [yr^{-1}]	15–25 jupiters, 2–3 earths	15–35 jupiters, 4–10 saturns
upper limit on planetary	4–7 % (jupiters)	3–7 % (jupiters)
abundance within 3 years	$\sim 40\%$ (earths)	10–30 % (saturns)
location of parent stars	Galactic disk and bulge	M31
extraction of		mostly difficult
planet parameters	fair in many cases	or even impossible

References

- Bennett, D. P., & Rhie, S. H. 1996, ApJ, 472, 660
Covone, G., de Ritis, R., Dominik, M., & Marino, A. A. 2000, A&A, 357, 816
Crofts, A. P. S. 1992, A&A, 399, L43
Dominik, M. 2002, in Napoli Series on Physics & Astrophysics, 6, General Relativity, Cosmology and Gravitational Lensing, ed. G. Marmo, C. Rubano & P. Scudellaro (Napoli: Bibliopolis), 87
Dominik, M., et al. 2002, P&SS, 50, 299
Gould, A., & Loeb, A. 1992, ApJ, 396, 104
Griest, K., & Safizadeh, N. 1998, ApJ, 500, 37
Jetzer, Ph. 1994, A&A, 286, 426